

Title

Perinatal Mortality in Nigeria: Do Place of Delivery and Delivery Assistants Matter?

Names and institutional affiliations of authors

Samuel Oji Oti^{a, b*} and Clifford Odimegwu^c

^a *School of Public Health, University of Witwatersrand, Johannesburg, South Africa*

^b *African Population and Health Research Center, Nairobi, Kenya*

^c *Programme in Demography and Population Studies, University of Witwatersrand, Johannesburg, South Africa*

*Corresponding author

Dr. Samuel Oti (MBBS, MSc.Med)

African Population and Health Research Center

Longonot Road, Upper Hill

P.O. Box 10787, GPO 00100,

Nairobi

Kenya

Tel: +254 20 2720400

Fax: +254 20 2720380

Email: soti@aphrc.org

Email addresses:

SO: soti@aphrc.org

CO: Clifford.Odimegwu@wits.ac.za

Abstract

Background: Studies in developed countries have shown that deliveries which occur at home and under the supervision of unskilled attendants present a high risk of perinatal death. However, it is unclear, especially in sub-Saharan Africa (SSA) where studies are limited, on precisely how these factors may influence perinatal mortality. This is particularly of interest considering that majority of births in SSA occur at home under the supervision of unskilled attendants and perinatal mortality in the sub-region is about five times that of the developed world. This study aims to determine the extent to which these factors influence perinatal mortality in Nigeria in order to inform maternal and child health policy decisions.

Methods: This study uses secondary data from the 2003 Nigerian Demographic and Health Survey. Logistic regression analysis was used to determine the association of place of delivery and delivery assistants with perinatal mortality while controlling for other known risk factors.

Results: Of the 5783 live singleton births that were analyzed, 194 were perinatal deaths giving an estimated perinatal mortality rate (PNMR) of 72.4 per 1000 live births. However, the results show that place of delivery and delivery assistants were not significantly associated with perinatal mortality in Nigeria.

Conclusions: In this study, place of delivery and delivery assistants are not critical predictors of perinatal mortality in Nigeria. However in view of the high PNMR found in this study, we recommend improved skilled attendance and institutional delivery rates as part of efforts for better maternal and child health in developing countries.

Background

Perinatal mortality –that is deaths in the first week of life including still births– is at an unacceptably high level in developing countries, especially those in sub-Saharan Africa (SSA)[1]. Recent estimates show that the perinatal mortality rate (PNMR) in developed regions of the world is about 10 per 1000 live births compared with 50 per 1000 live births in less developed regions of the world [1]. These figures are particularly troubling because the PNMR is known to be a key health status indicator of a community [2]. Specifically, it is an important indicator of the quality of obstetric and paediatric care available in any setting but at the same time it is a major contributor to overall childhood mortality [3].

Reducing childhood mortality levels by two-thirds by the year 2015 is one of the Millennium Development Goals (MDG-4) set by the United Nations [4]. Recent evidence suggests that neonatal mortality –death of a newborn within the first 4 weeks of life– contributes more than 50% of infant mortality globally. Furthermore, of all neonatal deaths, three-quarters occur in the perinatal period. Therefore, achieving MDG-4 will require a substantial reduction in perinatal deaths in high mortality countries [5, 6]. Unfortunately, despite evidence of gains in reducing childhood mortality globally, deaths in the perinatal period have shown the least progress and hence threaten to reverse these gains [1].

In 2000, the World Health Organisation (WHO) estimated the number of perinatal deaths globally to be greater than 6.3 million, with about 98% of these deaths occurring in low and middle-income countries [1, 6]. Nigeria is a low-income country that has experienced a surge in perinatal deaths over the past few decades. An earlier estimate of perinatal deaths in Nigeria was thought to be around 30 per 1000 live births in 1990 [7]. This figure almost tripled to 86 per 1000 live births by the year 2000 [1]. This high level of perinatal mortality in Nigeria means that the country may be unable to attain MDG-4 by the year 2015.

The level of perinatal mortality in a community is said to be associated with the proportion of births that take place at a health facility and the coverage of skilled attendance at birth. Countries where skilled attendance and institutional delivery rates are low usually have a high PNMR [6]. For instance, in Nigeria, only 35.2% of live births are delivered with the supervision of a skilled attendant and Nigeria has a very high PNMR of 86 per 1000 live births [1, 8].

Whereas in the United States (U.S.), 99.3% of live births are assisted by a skilled attendant [9] and the PNMR is only 7 per 1000 live births [1]. Therefore, improving the levels of skilled attendance at birth (delivery assistants) and increasing the numbers of births that occur in a health facility rather than at home (place of delivery) in countries where PNMR is high may lead to better perinatal health outcomes and ultimately reduce childhood mortality in line with the MDGs.

In the developed world, there is a sizable body of literature that has examined the roles of place of delivery and delivery assistants in determining perinatal mortality [10-14]. On one hand, it is quite clear in the literature that unsupervised deliveries and those supervised by unskilled persons carry a significantly high risk of perinatal mortality. On the other hand it remains unclear whether delivery in a hospital is in itself safer than delivery at home under the supervision of a skilled attendant [15]. In other words, the debate in the developed world is focussed on whether home deliveries are safer than institutional deliveries, considering that almost all births, whether occurring in a hospital or at home, are at least supervised by a skilled attendant.

However in the developing world where perinatal mortality rates remain high, most births occur at home and are supervised by unskilled persons such as untrained traditional birth attendants and relatives. Very few studies have attempted to investigate whether or not place of birth and delivery assistants are critical predictors of perinatal mortality in developing countries. Where available, these studies have often only examined the effect of place of delivery on perinatal mortality without considering the type of delivery assistant [16]. Other studies have looked at deliveries occurring at home and supervised by an unskilled person and compared the perinatal outcome with deliveries in a health facility [17, 18]. Such studies ignore the fact that even in developing countries some home deliveries are supervised by trained persons such as midwives or nurses. It therefore remains unclear in the context of developing countries what the precise roles of place of delivery and delivery assistants are in determining perinatal mortality. This study therefore aims to determine the extent to which place of delivery and delivery assistants predict perinatal mortality in the Nigerian context

Methodology

Data source

Data from the 2003 Nigerian Demographic and Health Survey (NDHS 2003) was used for this analysis. The complete methodology employed in the NDHS 2003 is described in the final survey report [8]. Briefly, the NDHS 2003 is the third national demographic and health survey (DHS) conducted in Nigeria by the National Population Commission (NPC) and Macro International Incorporated. It is based on a nationally representative sample of 7864 households. All women aged 15-49 years in these households were sampled to be individually interviewed. Hence, 7985 women were sampled, out of which 95% of them were successfully interviewed. The NDHS 2003 collected information on fertility levels and preferences, awareness and use of family planning methods, maternal and child health, breastfeeding practices, nutritional status of women and young children, childhood mortality, use of bed nets, female genital cutting, marriage, sexual activity, and awareness and behaviour regarding HIV/AIDS and other sexually transmitted infections in Nigeria.

Estimating the number of births

There were a total of 6029 observations in the dataset representing the total number of live births born to the interviewed mothers in the 5 years preceding the NDHS 2003. Out of this number, 246 live births were excluded because they were outcomes of multiple births. Therefore a total of 5783 live births were analyzed in this study, out of which 194 were deaths that occurred within the first seven days of life. These deaths do not include still births as this information was impossible to ascertain in the NDHS 2003.

Exposure and Outcome measures

The main explanatory variable is a composite of place of delivery and delivery assistants and is categorized as follows: “*Skilled institutional deliveries*”, “*Skilled home deliveries*”, “*Unskilled home deliveries*” and “*Unskilled institutional deliveries*”. We defined the term “skilled” as births supervised by doctors and or nurses/midwives;

while “unskilled” refers to births assisted by traditional birth attendants (TBA) and relatives/friends. The term “institutional” refers to births occurring in a health facility including public and private hospitals, clinics, health centers and health posts. While “home” refers to deliveries occurring outside a health facility, usually in a place of residence.

Other known controlling factors included in this study are maternal socio-economic and bio-demographic variables such as *Maternal age at birth*, *Highest level of education of mother*, *Region* (Geo-political location), *Place of residence* (rural/urban), *Religion*, *Wealth index*, *Maternal parity*, and number of *Antenatal visits in pregnancy*. Variables related to the child include *Birth size* (as reported subjectively by the mother) and *Gender of child*. Note that birth size is used as a proxy for birth weight since only about 10% of the respondents in the NDHS 2003 provided actual figures for birth weight. These factors have been shown in previous studies to be associated with perinatal mortality [12, 13, 16, 18-24].

The outcome variable used was the status of births by the first week of life –perinatal mortality. That is, each of the babies born to the interviewed women were categorized as alive or dead depending on whether or not the newborn survived the first seven days of life. Those babies who died within this period were compared to those who survived the same period. Note that this variable does not include information on still births which is not available in the primary dataset. Hence all analysis performed using this variable excludes still births.

Statistical Analysis

All data analysis was performed using STATA 10 statistical software. We used cross tabulations to describe and compare maternal and child characteristics by our main explanatory variable. This was followed by a bivariate analysis to examine the relationship between each variable of interest and the outcome variable –perinatal mortality (coded as zero if child survived the perinatal period and one if child died within this period). The final stage involved the use of stepwise logistic regression modeling to estimate the effect of place of delivery and delivery assistants on perinatal mortality, having controlled for the other potential risk factors.

Findings

The main objectives of this study were to estimate the level of perinatal mortality in Nigeria and to examine the influence of place of delivery and delivery assistants on perinatal mortality using data from the NDHS 2003. Out of 5783 singleton live births that occurred in the five years preceding the survey, there were 194 deaths before the 7th day of life giving an early neonatal mortality rate (ENMR) of 33.5 per 1000 live births. However, to determine the perinatal mortality rate (PNMR) we needed to estimate the still births rate (SBR), which as previously mentioned is unavailable in the primary data source. Hence, to estimate the SBR, we used formulae adopted by the WHO for countries where information on still births is lacking or inadequate [1].

The formula used is $SBR/ENMR=K$; Where K is a constant value unique to each of the six world regions: for Africa, WHO estimates K to be 1.2. The SBR is computed by multiplying the ENMR by K and therefore the SBR for Nigeria is 40.2 (33.5×1.2) per 1000 live births. The total number of still births (SBN) given the number of live births in the study population is 242 or $(40.2 \times 5783) / (1000 - 40.2)$. The total number of births (TNB) is given by (SBN + LB) and is therefore 6025 ($242 + 5783$). The PNMR which is given by $(SBN + LB) / (TNB)$ is equal to 72.4 per 1000 births, that is $(242 + 194) / 6025 \times 1000$.

Characteristics of births by place of delivery and delivery assistants

Unskilled home deliveries accounted for majority (60%) of the births in this study. Skilled institutional deliveries accounted for 34% while skilled home deliveries and unskilled institutional deliveries accounted for 4% and 2% of births respectively. Table 1 shows the distribution of selected maternal and child characteristics by place of delivery and delivery assistants. Unskilled home deliveries had the largest proportions of mothers living in rural areas (76%), who were poor (59%), who had no education (70.4%), and who had more than 5 previous live births (45%). Also, unskilled home deliveries had the lowest proportion of mothers who had 4 or more antenatal visits in pregnancy (28%). Skilled institutional deliveries had the highest proportion of births occurring in urban areas (53%), mothers who had 4 or more antenatal visits (85.6%), mothers who were educated above secondary level (54%), and also had the lowest proportion of mothers who were poor (22%).

Skilled home deliveries had the highest proportions of mothers aged 35 years or older at birth (29%) and the highest proportion of babies whose birth size was reported as below average or small by their mothers (20.4%). Unskilled

institutional deliveries had the largest proportion of mothers who had 2- 4 previous live births (56%) and the lowest proportion of babies whose birth size was reported as below average or small by their mothers (10.4%). All these differences in characteristics by place of delivery and delivery assistants were statistically significant. However, there was no statistical difference in the gender of the child by place of birth and delivery assistants.

Table 1. Maternal and child characteristics by place of delivery and delivery assistant.									
	Skilled institutional deliveries (n)	%	Skilled home deliveries (n)	%	Unskilled home deliveries (n)	%	Unskilled institutional deliveries (n)	%	p-value
Variables									
Maternal age at birth									<0.0001
<i><19 years</i>	89	4.6	25	9.9	259	7.5	4	5.2	
<i>20 - 34 years</i>	1380	71.4	154	61.1	2448	71.0	54	70.1	
<i>35+ years</i>	463	24.0	73	29.0	740	21.5	19	24.7	
	1932	100	252	100	3447	100	77	100	
Region									<0.0001
<i>North central</i>	425	22.0	47	18.7	480	13.9	15	19.5	
<i>North east</i>	245	12.7	66	26.2	1098	31.9	8	10.4	
<i>North west</i>	206	10.7	48	19.0	1466	42.5	6	7.8	
<i>South east</i>	356	18.4	28	11.1	86	2.5	5	6.5	
<i>South south</i>	273	14.1	35	13.9	201	5.8	26	33.8	
<i>South west</i>	427	22.1	28	11.1	116	3.4	17	22.1	
	1932	100	252	100	3447	100	77	100	
Place of residence									<0.0001
<i>Urban</i>	1031	53.4	106	42.1	828	24.0	31	40.3	
<i>Rural</i>	901	46.6	146	57.9	2619	76.0	46	59.7	
	1932	100	252	100	3447	100	77	100	
Highest educational level of Mother									
<i>No education</i>	345	17.9	94	37.3	2427	70.4	18	23.4	
<i>primary</i>	551	28.5	84	33.3	711	20.6	32	41.6	
<i>secondary</i>	852	44.1	72	28.6	294	8.5	24	31.2	
<i>higher</i>	184	9.5	2	0.8	15	0.4	3	3.9	

	1932	100	252	100	3447	100	77	100	
Marital status									<0.0001
<i>Never married</i>	73	3.8	2	0.8	30	0.9	2	2.6	
<i>Currently married</i>	1780	92.1	241	95.6	3306	95.9	72	93.5	
<i>Formerly married</i>	79	4.1	9	3.6	111	3.2	3	3.9	
	1932	100	252	100	3447	100	77	100	
Gender of child									0.16
<i>Male</i>	1002	51.9	127	50.4	1721	49.9	47	61.0	
<i>Female</i>	930	48.1	125	49.6	1726	50.1	30	39.0	
	1932	100	252	100	3447	100	77	100	
Wealth Index									<0.0001
<i>Poor</i>	417	21.6	72	28.6	2049	59.4	27	35.1	
<i>Middle</i>	800	41.4	131	52.0	1252	36.3	33	42.9	
<i>Rich</i>	715	37.0	49	19.4	146	4.2	17	22.1	
	1932	100	252	100	3447	100	77	100	
Religion									<0.0001
<i>Christian</i>	1276	66.0	118	46.8	734	21.3	49	63.6	
<i>Islam</i>	637	33.0	131	52.0	2621	76.1	27	35.1	
<i>Traditional /other</i>	19	1.0	3	1.2	87	2.5	1	1.3	
	1932	100	252	100	3442	100	77	100	
Maternal parity									<0.0001
<i>primiparous</i>	348	18.0	45	17.9	330	9.6	9	11.7	
<i>2 – 4 births</i>	980	50.7	102	40.5	1558	45.2	43	55.8	
<i>≥ 5 births</i>	604	31.3	105	41.7	1559	45.2	25	32.5	
	1932	100	252	100	3447	100	77	100	
Antenatal visits in pregnancy									<0.0001
<i>no visit</i>	36	2.8	24	14.5	1161	55.3	15	30	
<i>less than 4</i>	146	11.5	25	15.2	347	16.5	5	10	
<i>4 or more</i>	1086	85.6	116	70.3	593	28.2	30	60	
	1268	100	165	100	2101	100	50	100	

Birth size									<0.0001
<i>Average</i>	811	42.2	108	43.2	1500	43.9	33	42.9	
<i>Larger than average</i>	880	45.8	91	36.4	1392	40.8	36	46.8	
<i>Smaller than average</i>	232	12.1	51	20.4	522	15.3	8	10.4	
	1923	100	250	100	3414	100	77	100	

*Chi-square p-value

Table 2 shows that neither unskilled home deliveries (unadjusted OR 1.04, 95% CI 0.73 - 1.48) nor unskilled institutional deliveries (unadjusted OR 1.68, 95% CI 0.62 - 4.59) were significantly associated with a higher risk of perinatal deaths in comparison to skilled institutional deliveries. Also, skilled home deliveries did not have a significantly higher risk of perinatal deaths in comparison to skilled institutional deliveries (unadjusted OR 1.14, 95% CI 0.52 - 2.49). However, births in rural areas (unadjusted OR 1.57, 95% CI 1.07 - 2.32), babies whose birth size was smaller than average (unadjusted OR 1.55, 95% CI 1.05 - 2.31) and those whose mothers practiced “traditional/other” religion (unadjusted OR 5.12, 95% CI 3.14 - 8.33) were all significantly associated with an increased risk of perinatal deaths. On the other hand, female babies (unadjusted OR 0.56, 95% CI 0.41 - 0.77) and those born to “rich” mothers had a significantly lower risk of perinatal deaths (unadjusted OR 0.43, 95% CI 0.25 - 0.75).

Variables	n†	N‡	%	Crude odds ratio (95% CI)	P-value
Place of delivery and delivery assistants					
<i>Skilled institutional deliveries</i>	61	1932	3.2	1	
<i>Skilled home deliveries</i>	9	252	3.6	1.13 (0.52 - 2.49)	0.75
<i>Unskilled home deliveries</i>	113	3447	3.3	1.04 (0.73 - 1.48)	0.83
<i>Unskilled institutional deliveries</i>	4	77	5.2	1.68 (0.62 - 4.59)	0.31
Maternal age at birth					
<i><19 years</i>	19	381	5.0	1	
<i>20 - 34 years</i>	128	4084	3.1	0.62 (0.35 - 1.07)	0.09
<i>35+ years</i>	47	1318	3.6	0.71 (0.39 - 1.25)	0.24

Place of residence					
<i>Urban</i>	50	2024	2.5	1	
<i>Rural</i>	144	3759	3.8	1.57 (1.07 - 2.32)	0.022
Gender of child					
<i>Male</i>	125	2940	4.3	1	
<i>Female</i>	69	2843	2.4	0.56 (0.41 - 0.77)	<0.0001
Wealth Index					
<i>Poor</i>	108	2614	4.1	1	
<i>Middle</i>	69	2239	3.1	0.74 (0.53 - 1.03)	0.07
<i>Rich</i>	17	930	1.8	0.43 (0.25 - 0.75)	0.003
Religion					
<i>Christian</i>	74	2212	3.3		
<i>Islam</i>	103	3453	3.0	0.89 (0.63 - 1.25)	0.5
<i>Traditional/other</i>	17	113	15.0	5.12 (3.14 - 8.33)	<0.0001
Maternal parity					
<i>primiparous</i>	31	734	4.2	1	
<i>2 – 4 births</i>	77	2712	2.8	0.66 (0.43 - 1.02)	0.06
<i>≥5 births</i>	86	2337	3.7	0.87 (0.57 - 1.31)	0.49
Birth size					
<i>Average</i>	81	2458	3.3	1	
<i>Larger than average</i>	61	2412	2.5	0.76 (0.56 - 1.03)	0.08
<i>Smaller than average</i>	41	815	5.0	1.55 (1.05 - 2.31)	0.03

† Number of deaths within the first seven days of life.

‡ Total number of live births

In the multivariate analysis shown in table 3, we fit our main explanatory variable representing place of delivery and delivery assistants into a final model while controlling for the risk factors that were significantly associated ($p < 0.05$) with perinatal mortality in the unadjusted analysis. Place of delivery and delivery assistants remained non-significant in this model while births in rural areas, female babies, being smaller than average in size at birth, and being born to a mother who practiced “traditional/other” religion remained significantly associated with perinatal mortality. However, being born to a rich mother was no longer a significant risk factor in the adjusted model.

Table 3. Multivariate analysis for perinatal mortality.

Variables	Adjusted odds ratio (95% CI)	P-value
Place of delivery and delivery assistants		
<i>Skilled institutional deliveries</i>	1	
<i>Skilled home deliveries</i>	1.01 (0.44 - 2.26)	0.99
<i>Unskilled home deliveries</i>	0.8 (0.52 - 1.21)	0.29
<i>Unskilled institutional deliveries</i>	1.51 (0.53 - 4.32)	0.44
Place of residence		
<i>Urban</i>	1	
<i>Rural</i>	1.49 (1.02 - 2.18)	0.041
Gender of child		
<i>Male</i>	1	
<i>Female</i>	0.57 (0.42 - 0.78)	<0.0001
Wealth Index		
<i>Poor</i>	1	
<i>Middle</i>	0.89 (0.63 - 1.27)	0.53
<i>Rich</i>	0.58 (0.31 - 1.08)	0.1
Religion		
<i>Christian</i>	1	
<i>Islam</i>	0.93 (0.61 - 1.41)	0.74
<i>Traditional/other</i>	4.09 (2.17 - 7.72)	<0.0001
Birth size		
<i>Average</i>	1	
<i>Larger than average</i>	0.76 (0.56 - 1.03)	
<i>Smaller than average</i>	1.56 (1.05 - 2.33)	0.03

Discussion

This study purposed to estimate the level of perinatal mortality in Nigeria and to examine the role of place of delivery and delivery assistants in determining perinatal mortality, while taking other selected risk factors into consideration. Using the NDHS 2003 as our data source, we obtained an ENMR of 33.5 per 1000 live births and an estimated PNMR of 72.4 per 1000 live births. We did not find any significant association between our main explanatory variable –place of delivery and delivery assistants– with perinatal mortality. However, other factors such as being a female child, religion of the mother, size of the child at birth and type of place of residence (rural versus urban) were found to be significantly associated with perinatal mortality.

The lack of association between place of delivery and delivery assistants with perinatal mortality is inconsistent with previous studies [10, 12, 18] and is especially surprising for a developing country like Nigeria where most deliveries occur at home and are not supervised by a skilled attendant [25]. In the developed world where most deliveries occur at a health facility and are attended by skilled health personnel, the above results will not be unusual [14]. One possible explanation for this result may be that majority of those who deliver at a health facility only do so if there are complications in labour by which time the health of the fetus has been seriously compromised. If this is combined with a lack of high quality perinatal care units as is the case in many health facilities in Nigeria, the result may be a level of perinatal mortality in skilled institutional deliveries that is not very different from unskilled home delivery rates.

Another explanation may be that perhaps a good number of the “unskilled” persons such as TBA who attend to majority of births in Nigeria are in fact really good at what they do and have succeeded in avoiding significant differences in their perinatal death rates. Some previous studies in Nigeria revealed that one of the reasons why women in Nigeria patronized TBA was simply due to their positive experiences –they have had successful deliveries in the hands of TBA [26, 27]. However, these studies emphasize that the training, retraining and monitoring of TBA is crucial to ensure that the quality of services they offer are at an acceptable standard of safe birth practices.

We however found some other factors that were significantly associated with perinatal mortality. Firstly, we found that female children had a lower risk of dying in the perinatal period compared to males. It has been suggested that male children are more prone to perinatal deaths than females for biological reasons [24]. By contrast, some other studies have found a higher risk of perinatal deaths in females than males. However, these studies justified their findings by arguing that gender preference was practiced in their study populations [28, 29]. There is no evidence to suggest that gender preference is practiced in Nigeria at a general level. Thus we can conclude that natural selection which gives female children the survival edge over males at this early stage of life is what prevails in the Nigerian context.

We also found that being born to a mother who practiced “traditional/other” religion was found to be significantly associated with an increased risk of perinatal death. The influence of religion on perinatal mortality may be linked to the beliefs and myths they may have concerning child birth. This was the case with the Faith Assembly people in Indiana, U.S., who believed that child bearing was an act of God not to be interfered with [12]. Unfortunately, this group had a perinatal mortality rate that was thrice that of the general population. We explored this association further by introducing an interaction term between religion and gender of the child on our final model (results not shown). We found that female children who were born to mothers who practiced “traditional/other” religion had a higher risk (OR 4.17 95% CI 1.83 - 9.51) of perinatal deaths compared to male children born to “Christian” mothers. It is difficult to say whether or not this finding reflects gender preference among the subgroup of traditionalists and further qualitative research may be useful.

Children who were reported to be “smaller than average” in size at birth, were also found to have a significantly higher risk of perinatal deaths than those who were average in size. This may reflect the increased susceptibility of low birth weight babies to infections and other abnormalities [19, 30]. The prevailing argument is that low birth weight is a frequent intermediate factor linking other risk factors such as socioeconomic status to perinatal mortality [20, 30]. While this may be true in our study as well, caution must be exercised because we used a subjective measure of birth size as a proxy for birth weight.

Finally, we also found that rural dwellers had a higher risk of perinatal deaths when compared to urban residents. This is supported by several studies [21, 22, 30], which generally attribute this finding to better access to perinatal care facilities in the urban areas reflecting social inequalities between the two groups.

Limitations

The quality of any DHS data depends on the completeness with which births and deaths are reported. Deaths in the first week of life are often anonymous in that the new born die and are quickly buried before they are even named. This often blurs information on these deaths and the circumstances surrounding some of them. Hence the possibility of selective under-reporting of perinatal deaths is a real threat to the data quality of this study. In order to assess if there has been selective under-reporting, the DHS uses the ratio of deaths under 7 days to all neonatal deaths as an indicator. If this ratio (expressed as a percentage) is unusually low, then it is very likely that selective under-reporting of perinatal deaths has occurred. In the case of the NDHS 2003, this ratio was quite high —74% for the period 0-4 years preceding the study [8]. Hence we can conclude that although selective under-reporting of perinatal deaths was a potential limitation of our study, this was not the found to be the case.

Another major limitation of our study is the issue of uncontrolled confounding. For example, pregnancy and labour complications are a strong confounder for perinatal deaths [31]. Home deliveries and those assisted by unskilled persons are associated with a high risk of labour complications. And labour complications, which are a risk with every pregnancy, may be associated with increased risk of perinatal death. Some studies categorize each pregnancy by risk status and have found associations with place of delivery and perinatal mortality among high risk pregnancies but not necessarily low risk pregnancies [18]. However, full and accurate accounts of pregnancy and or labour complications are not available in the NDHS 2003 and therefore this potential confounder could not be included in our analysis.

Also, we found that information in our data source on still births is unclear –no distinction was made in the primary data source as to whether those pregnancies which did not result in live births were due to abortions, miscarriages or still births. Therefore the SBR in this study was estimated using a methodology developed by the WHO. This

involved the use of regional estimates of the SBR/PNMR ratio to calculate the SBR in countries where no data on still births are available [1]. While the WHO assures that every effort has been made to arrive at reliable and balanced estimates, it also reiterates that it is not possible to calculate precise mortality rates, because reliable population-based data are not available for all countries and for all years. Nevertheless, the WHO emphasizes that regional estimates should be regarded as the best estimates possible on the basis of the information available.

Conclusion

Our study did not find any effect of place of delivery and delivery assistants on perinatal mortality. This lack of association may be due to the limitations of our study previously described. However, we found a PNMR in our study that is comparable to existing estimates and which by every standard is very high.

We therefore recommend that as far as health policy and planning is concerned, the Nigerian government and its development partners need to do more in order to attain significant and sustained improvement in maternal and child health and especially perinatal health. One of the United Nations MDG indicators in this regard is the proportion of deliveries supervised by a skilled attendant and we believe that the reported proportion of less than 40% for skilled delivery rates in Nigeria is unacceptably low. However, skilled delivery rates and institutional delivery rates go hand in hand. Health workers can not be trained and retained in a vacuum. Therefore, improved skilled and institutional delivery rates can be increased via building and equipping accessible primary health care facilities and training more health personnel to attend to these facilities especially in the rural areas. Retaining a skilled health workforce especially in rural areas is a challenge for many developing countries. Hence, incentives to motivate and retain rural health workers may be necessary. These incentives may include but are not limited to financial rewards, availability of career development opportunities, access to continuing education, and service recognition/appreciation.

Although reducing perinatal mortality is only one aspect in the drive to reduce childhood mortality by two-thirds in 2015 in line with MDG-4, it will be an essential step which must be taken if sustainable progress is to be made as regards child health.

Acknowledgements

We wish to acknowledge Macro International Inc., for granting us permission to use data from the 2003 Nigerian Demographic and Health Survey. We also acknowledge the contribution of Dr. Catherine Kyobutungi and Dr. Latifat Ibisomi who critically reviewed this paper for important intellectual content.

Competing interest

The authors declare that they have no competing interests.

Ethical Approval

This study involved secondary analysis of a public domain data set –NDHS 2003. Permission to use this data set was obtained in writing from Macro International Incorporation via the DHS website (www.measuredhs.com). The interview process via which the survey was conducted included the documentation of informed consent which was given by the respondents [8]. Respondent confidentiality is in tact as no names are included in the data set and therefore the respondents can not be traced by the researcher. Finally, the research protocol was submitted for review by Ethics Committee of the University of the Witwatersrand, Johannesburg –South Africa, and ethical clearance was obtained before the research began.

Authors' contributions

SO did the data analysis, literature review and wrote the first draft of the manuscript and contributed to the interpretation of the findings.

CO conceptualized this study and contributed to the writing of the paper and interpretation of the findings.

Both authors read and approved the final manuscript

References

1. WHO. World Health Report. Neonatal & Perinatal Mortality:Country, Regional and Global estimates. Geneva; 2006.
2. Wilkinson D. Reducing perinatal mortality in developing countries. Health policy and planning. 1997; 12(2):160-5.
3. Richardus J, Graafmansp W, Verloove-Vanhorick S, Mackenbach J. The Perinatal Mortality Rate as an Indicator of Quality of Care in International Comparisons. Medical Care. 1998;36(1):54-66.
4. The United Nations. The Millenium Development Goals Report. New York; 2006.
5. Jose M, Paul I, Z, B., Marjorie K, *et al.* Neonatal survival:a call for action. Lancet 2005;365:1189-97.
6. Lawn J, Cousens S, Zupan J. 4 million deaths. When? Where? Why? . Lancet 2005;365:891-900.
7. WHO. Perinatal mortality:a listing of available information. Geneva; 1996.
8. National Population Commission [Nigeria]., ORC Macro. Nigeria Demographic and Health Survey 2003: The Final Report. Calverton, Maryland, USA; 2004.
9. WHO. Health situation in the Americas. Basic indicators. Washington DC; 2006.
10. Burnett C, Jones J, Rooks J, Chen C, Tyler CJ, Miller C. Home delivery and neonatal mortality in North Carolina. JAMA 1980 244(24):2741-5.
11. Campbell R, Macdonald-Davies I, Macfarlane A, Beral V. Home births in England and Wales,1979: perinatal mortality according to intended place of delivery. BMJ 1984 289:721-4.
12. Kaunitz M, Spence C, Danilson T, Rochat R, Grimas D. Perinatal mortality in a religious group avoiding obstetric care. American Journal of Obstetrics and Gynecology. 1984;150:826-30
13. Parazzini F, LaVecchia C. Perinatal mortality and place of birth in Italy. Am J of Public Health. 1988;78(6):706-7.
14. Schramm W, Barnes D, Bakewell J. Neonatal mortality in Missouri home births, 1978-84. Am J of Public Health. 1987;77(8):930-5.
15. Durand AM. The safety of home births: The farm study. Am J of Public Health 1992 82(3):450 – 3.
16. Mcdermott J, Steketee R, Wirima J. Perinatal mortality in rural Malawi. Bulletin of the World Health Organization. 1996;74 (2):165-71.
17. Voorhoeve A, Kars C, Ginneken J. Modern and traditional antenatal and delivery care. In: van Ginneken JK, Muller AS, editors. Maternal and Child health in Rural Kenya. Kent: Croom Helm 1985. p. 309-22.
18. Walraven G, Mkanje R, Roosmalen J, Van Dongen P, Dolmas W. Perinatal mortality in home births in rural Tanzania. European Journal of Obstetrics and Gynecology and Reproductive Biology 1995;58:131-4.
19. Borrell C, Grera E, Ricart M, Pasarin M, Salvador J. Social Inequalities in Perinatal Mortality in a Southern European City. European Journal of Epidemiology 2003 18(1):5-13.
20. Forsass E, Gissler M, Sihvonen M, Hemminki E. Maternal predictors of perinatal mortality: the role of birthweight. Int J Epid 1999 28:475-8.
21. Golding J, Greenwood R, McCaw-Binns A, Thomas P. Associations between social and environmental factors and perinatal mortality in Jamaica. Paediatr Perinat Epidemiol. 1994;8(1):17-39.
22. Habib N, Lie R, Oneko O, Shao J, Bergsjø P, Daltveit A. Socio demographic characteristics and perinatal mortality among singletons in North East Tanzania: a registry-based study J Epidemiol Community Health 2008;62:960-5.
23. Shah N, Shah M, Khalaf A, Mustafa M, Al-Sayed. A. Searching for socioeconomic risk factors in perinatal mortality in Kuwait: a case-control study. Soc Sci & Med. 2000;51:539-50.
24. Wessel H, Cnattingius S., Dupret A., Reitmaier P., S. B. Risk factors for perinatal deaths in Cape Verde. Paediatric Perinatal Epidemiology 1998;12 (1):25–36.
25. WHO. Proportion of births attended by a skilled attendant -2007 updates. Geneva: Department of Reproductive Health and research, WHO; 2007 [cited 2007 August 21]; Available from: www.who.int/reproductive-health.
26. Imogie A, Agwubike E, Aluko K. Assessing the role of Traditional Birth Attendants in Health Care delivery in Edo State, Nigeria Afr J Rep Health. 2002 6(2):94-100.

27. Salako A, Daniel O. Identifying the training needs of Traditional Birth Attendants. *Tropical Doctor*. 2007 37:6-10.
28. Nielsen B, Liljestrand J, Hedegaard M, Thilsted S, Joseph A. Reproductive pattern, perinatal mortality, and sex preference in rural Tamil Nadu, south India: community based, cross sectional study. *BMJ*. 1997;314(7093):1521-4.
29. Wu Z, Viisainen K, Wang Y, Hemminki E. Perinatal mortality in rural China: a retrospective cohort. *BMJ* 2003;327:1319-22.
30. Silins J, Semenciw M, Morrison H, *et al*. Risk factors for perinatal mortality in Canada. *Canadian Medical Association Journal* 1985;133(12):1214-9.
31. Weiner R, Ronsmans C, Dorman E, Jilo H, Muhoro A, C. S. Labour complications remain the most important risk factors for perinatal mortality in rural Kenya. *Bulletin of the World Health Organization*. 2003;81(7).

